

# Physics Reference Tables

## PHYSICAL CONSTANTS AND CONVERSION FACTORS

Acceleration due to gravity	$g$	9.8 m/s/s or m/s <sup>2</sup>
Speed of light in a vacuum	$c$	$3.00 \times 10^8$ m/s
Electron rest mass	$m_e$	$9.11 \times 10^{-31}$ kg
Electron charge	$e$	$1.6 \times 10^{-19}$ C
Proton rest mass	$m_p$	$1.67 \times 10^{-27}$ kg
Gravitation constant	$G$	$6.67 \times 10^{-11}$ N•m <sup>2</sup> /kg <sup>2</sup>
Coulomb's law constant	$k$	$9.0 \times 10^9$ N•m <sup>2</sup> /C <sup>2</sup>
Speed of sound at STP		331 m/s

## THE INDEX OF REFRACTION FOR COMMON SUBSTANCES

( $\lambda = 5.9 \times 10^{-7}$  m)

Air	1.00
Alcohol	1.36
Canada Balsam	1.53
Corn Oil	1.47
Diamond	2.42
Glass, Crown	1.52
Glass, Flint	1.61
Glycerol	1.47
Lucite	1.50
Quartz, Fused	1.46
Water	1.33

## WAVELENGTHS OF LIGHT IN A VACUUM

Violet	$4.0 - 4.2 \times 10^{-7}$ m
Blue	$4.2 - 4.9 \times 10^{-7}$ m
Green	$4.9 - 5.7 \times 10^{-7}$ m
Yellow	$5.7 - 5.9 \times 10^{-7}$ m
Orange	$5.9 - 6.5 \times 10^{-7}$ m
Red	$6.5 - 7.0 \times 10^{-7}$ m

## HEAT CONSTANTS

	Specific Heat (average) (kJ/kg•°C) or (J/g•°C)	Melting Point (°C)	Boiling Point (°C)
Alcohol (ethyl)	2.43 (liq.)	-117	79
Aluminum	0.90 (sol.)	660	2467
Ammonia	4.71 (liq.)	-78	-33
Copper	0.39 (sol.)	1083	2567
Iron	0.45 (sol.)	1535	2750
Lead	0.13 (sol.)	328	1740
Mercury	0.14 (liq.)	-39	357
Platinum	0.13 (sol.)	1772	3827
Silver	0.24 (sol.)	962	2212
Tungsten	0.13 (sol.)	3410	5660
Water (solid)	2.05 (sol.)	0	—
Water (liquid)	4.18 (liq.)	—	100
Water (vapor)	2.01 (gas)	—	—
Zinc	0.39 (sol.)	420	907

# FORMULAS

## MECHANICS

$$\begin{aligned}\bar{v} &= \frac{\Delta s}{\Delta t} \\ \bar{v} &= \frac{v_f + v_i}{2} \\ \bar{a} &= \frac{\Delta v}{\Delta t} \\ \Delta s &= v_i \Delta t + \frac{1}{2} a (\Delta t)^2 \\ v_f^2 &= v_i^2 + 2a \Delta s \\ F &= ma \\ w &= mg \\ F &= \frac{G m_1 m_2}{r^2} \\ p &= mv \\ F \Delta t &= \Delta(mv) \\ \tau &= F d \sin \theta \\ \sum \tau_{cw} - \sum \tau_{ccw} &= 0 \\ a_c &= \frac{v^2}{r} \\ F_c &= \frac{mv^2}{r}\end{aligned}$$

$a$  = acceleration  
 $a_c$  = centripetal acceleration  
 $v$  = velocity  
 $r$  = radius  
 $F$  = force  
 $F_c$  = centripetal force  
 $\theta$  = angle  
 $g$  = acceleration due to gravity  
 $G$  = universal gravitational constant  
 $m$  = mass  
 $p$  = momentum  
 $\Delta s$  = displacement  
 $t$  = time  
 $\tau$  = torque  
 $w$  = weight

## ELECTRICITY AND MAGNETISM

$$\begin{aligned}F &= \frac{k q_1 q_2}{r^2} & r &= \text{distance between centers} \\ V &= \frac{W}{q} & F &= \text{force} \\ I &= \frac{\Delta q}{\Delta t} & I &= \text{current} \\ I &= \frac{V}{R} & k &= \text{electrostatic constant} \\ P &= VI & P &= \text{power} \\ F &= qvB & q &= \text{charge} \\ V &= Blv & R &= \text{resistance} \\ W &= Pt = VI t & V &= \text{electrical potential difference} \\ & & W &= \text{energy} \\ & & B &= \text{flux density} \\ & & \ell &= \text{length of a conductor} \\ & & v &= \text{velocity}\end{aligned}$$

### Series Circuits

$$\begin{aligned}I_t &= I_1 = I_2 = I_3 = \dots \\ V_t &= V_1 + V_2 + V_3 + \dots \\ R_t &= R_1 + R_2 + R_3 + \dots\end{aligned}$$

### Parallel Circuits

$$\begin{aligned}I_t &= I_1 + I_2 + I_3 + \dots \\ V_t &= V_1 = V_2 = V_3 = \dots \\ \frac{1}{R_t} &= \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots\end{aligned}$$

# FORMULAS

## ENERGY

$$\begin{aligned}W &= F\Delta s \\P &= \frac{W}{\Delta t} = \frac{F\Delta s}{\Delta t} = F\bar{v} \\ \Delta PE_g &= mg\Delta h \\ KE &= \frac{1}{2}mv^2 \\ F &= kx \\ PE_s &= \frac{1}{2}kx^2 \\ \mu_s &= \frac{F_{s,\max}}{F_n}\end{aligned}$$

$F$  = force  
 $g$  = acceleration due to gravity  
 $h$  = height  
 $k$  = spring constant  
 $x$  = change in length of a spring from the equilibrium position  
 $KE$  = kinetic energy  
 $m$  = mass  
 $P$  = power  
 $PE_g$  = gravitational potential energy  
 $PE_s$  = potential energy stored in a spring  
 $\Delta s$  = displacement  
 $t$  = time  
 $v$  = velocity  
 $W$  = work  
 $\mu_s$  = coefficient of static friction

## INTERNAL ENERGY

$$\begin{aligned}Q &= mC_p\Delta T \\ Q &= \Delta E - W = (E_f - E_i) - W\end{aligned}$$

$Q$  = amount of heat  
 $C_p$  = specific heat  
 $T$  = temperature  
 $W$  = work  
 $E_f$  = final energy of the system  
 $E_i$  = initial energy of the system

## WAVE PHENOMENA

$$\begin{aligned}T &= \frac{1}{f} & c &= \text{speed of light in a vacuum} \\ v &= f\lambda & f &= \text{frequency} \\ n &= \frac{c}{v} & n &= \text{index of absolute refraction} \\ \sin \theta_{c\ air} &= \frac{1}{n} & T &= \text{period} \\ \frac{n_1}{n_2} &= \frac{\sin \theta_2}{\sin \theta_1} & v &= \text{speed} \\ \frac{n_1}{n_2} &= \frac{v_2}{v_1} & \lambda &= \text{wavelength} \\ && \theta &= \text{angle} \\ && \theta_{c\ air} &= \text{critical angle of incidence relative to air}\end{aligned}$$

## GEOMETRIC OPTICS

$$\begin{aligned}\frac{1}{d_o} + \frac{1}{d_i} &= \frac{1}{f} & d_i &= \text{image distance} \\ \frac{S_o}{S_i} &= \frac{d_o}{d_i} & d_o &= \text{object distance} \\ && S_i &= \text{image size} \\ && S_o &= \text{object size} \\ && f &= \text{focal length}\end{aligned}$$

## INVESTIGATIONS

$$\% \text{ Error} = \frac{\text{Accepted value} - \text{Experimental value}}{\text{Accepted value}} \times 100$$

# Electromagnetic Spectrum

